

Disclaimer

This English translation is prepared by Marketing Authorization Holder.

The Japanese language package insert is the official document, and this translation is provided solely as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.

The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.

This translation shall not be reproduced, reprinted or used for profit.

Since this translation is prepared in July 2025 based on the Japanese text current at that time, the translation may not reflect the latest information, due to continuous revision of package inserts.

The latest Japanese text is available on PMDA website.

Kracie

Oct 2023 (1st version)

Standard Commodity Classification No. of Japan	
875200	
Approval No.	16100AMZ03513000
Date of Initial Marketing in Japan	Oct 1986

Kampo product Kracie **Kakkonto** Extract Fine Granules

Storage : Store at room temperature
Shelf Life : 3 years

KB-1

EK-1

3. COMPOSITION AND PRODUCT DESCRIPTION

3.1 Composition

Brand name	Kracie Kakkonto Extract Fine Granules
Active ingredients	The daily dose of this product (7.5 g) contains 5,200 mg of JP Kakkonto extract, derived from a mixture of the following crude drugs.
	JP Pueraria Root 8.0 g
	JP Jujube 4.0 g
	JP Ephedra Herb 4.0 g
	JP Glycyrrhiza 2.0 g
	JP Cinnamon Bark 3.0 g
	JP Peony Root 3.0 g
JP Ginger 1.0 g	
Inactive ingredients	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide

(JP: The Japanese Pharmacopoeia)

3.2 Product Description

Dosage form	Fine granules
Color	Light red-brown to brown
Odor	Characteristic odor
Taste	Slightly bitter and sweet
ID code	KB-1 (3.75 g per sachet)
	EK-1 (2.5 g per sachet)

4. INDICATIONS

Common cold, head cold, headache, shoulder muscle stiffness, myalgia, hand and shoulder aches

6. DOSAGE AND ADMINISTRATION

The usual adult dosage for oral use is 7.5 g daily in two or three divided doses before or between meals.

The dosage may be adjusted according to the patient's age, body weight, and symptoms.

8. IMPORTANT PRECAUTIONS

8.1 When this product is used, the patient's "SHO" (constitution/symptoms) should be taken into consideration. The patient's progress should be carefully monitored, and if no improvement in symptoms or findings is observed, continuous administration should be avoided.

8.2 Since this product contains Glycyrrhiza, careful attention should be paid to the serum potassium level, blood pressure, etc. [See Sections 10.2, 11.1.1, 11.1.2.]

8.3 When this product is used in combination with other Kampo products, etc., attention should be paid to the duplication of the contained crude drugs.

9. PRECAUTIONS CONCERNING PATIENTS WITH SPECIFIC BACKGROUNDS

9.1 Patients with Complication or History of Diseases, etc.

9.1.1 Patients in a period of weakness after disease or with extremely weakened constitution

Adverse reactions are likely to occur, and the symptoms may be aggravated.

9.1.2 Patients with an extremely weak gastrointestinal tract

Anorexia, epigastric distress, nausea, vomiting, etc. may occur.

9.1.3 Patients with anorexia, nausea, or vomiting

These symptoms may be aggravated.

9.1.4 Patients with a significant sweating tendency

Excessive sweating, systemic weakness, etc. may occur.

9.1.5 Patients with cardiovascular disorders, including angina pectoris or myocardial infarction, or patients with a history of such disorders

The disease and its symptoms may be aggravated.

9.1.6 Patients with severe hypertension

The disease and its symptoms may be aggravated.

9.1.7 Patients with urination impaired

The disease and its symptoms may be aggravated.

9.1.8 Patients with hyperthyroidism

The disease and its symptoms may be aggravated.

9.2 Patients with Renal Impairment

9.2.1 Patients with severe renal disorder

The disease and its symptoms may be aggravated.

9.5 Pregnant Women

This product should be used in pregnant women or women who may possibly be pregnant only if the expected therapeutic benefits outweigh the possible risks associated with treatment.

9.6 Breast-feeding Women

Considering the therapeutic benefits and the benefits of breastfeeding, continuation or discontinuation of breastfeeding should be considered.

9.7 Pediatric Use

No clinical studies have been conducted in children.

9.8 Geriatric Use

Since the physiological functions are generally decreased in elderly patients, careful supervision is recommended; measures such as reducing the dose may be considered.

10. INTERACTIONS

10.2 Precautions for Co-administration (This drug should be administered with caution when co-administered with the following.)

Drugs	Signs, Symptoms, and Treatment	Mechanism and Risk Factors
Ephedra Herb-containing preparations Shoseiryuto Maoto Maobushisaishinto, etc. Ephedrine-containing preparations Ephedrine Hydrochloride dl-Methyl Ephedrine Hydrochloride Fexofenadine Hydrochloride/Pseudoephedrine Hydrochloride, etc. Monoamine Oxidase (MAO) inhibitors Selegiline Hydrochloride Rasagiline Mesilate, etc. Thyroid gland preparations Thyroxine Liothyronine, etc. Catecholamine preparations Adrenaline Isoprenaline, etc. Xanthine preparations Theophylline Diprophylline, etc.	Since insomnia, excessive sweating, tachycardia, palpitation, systemic weakness, mental excitement, etc. are likely to occur, this product should be administered with care by reducing the dosage, etc.	The sympathomimetic effect may be enhanced.
Glycyrrhiza-containing preparations Shakuyakukanzoto Hochuekkito Yokukansan, etc. Preparations containing glycyrrhizic acid and its salts Monoammonium Glycyrrhizinate/Glycine/L-cysteine Monoammonium Glycyrrhizinate/Glycine/DL-Methionine combination tablets, etc. [See Sections 8.2, 11.1.1, 11.1.2]	Pseudoaldosteronism is likely to occur. As a result of hypokalaemia, myopathy is likely to occur.	Since glycyrrhizic acid has an accelerating effect on potassium excretion in the renal tubules, an acceleration of decrease in the serum potassium level has been suggested.

11. ADVERSE REACTIONS

The following adverse reactions may occur. Patients should be carefully monitored, and if any abnormalities are observed, appropriate measures such as discontinuation of administration should be taken.

11.1 Clinically Significant Adverse Reactions

11.1.1 Pseudoaldosteronism (frequency unknown)

Pseudoaldosteronism such as hypokalaemia, blood pressure increased, retention of sodium/body fluid, edema, and body weight gain may occur. Patients should be carefully monitored (e.g., measurement of serum potassium levels), and if any abnormalities are observed, administration should be discontinued, and appropriate measures such as administration of potassium preparations should be taken. [See Sections 8.2, 10.2]

11.1.2 Myopathy (frequency unknown)

Myopathy may occur as a result of hypokalaemia. Patients should be carefully monitored, and if any abnormalities such as feelings of weakness, muscle cramp in extremities, or paralysis are observed, administration should be discontinued and appropriate measures such as administration of potassium preparations should be taken. [See Sections 8.2, 10.2]

11.1.3 Hepatic Impairment, jaundice (frequency unknown)

Hepatic impairment and/or jaundice with marked elevations of AST, ALT, Al-P, γ -GTP, etc. may occur.

11.2 Other Adverse Reactions

	Frequency unknown
Hypersensitivity	Rash, Redness, Pruritus, etc.
Autonomic	Insomnia, Excessive sweating, Tachycardia, Palpitations, Systemic weakness, Mental excitement, etc.
Gastrointestinal	Anorexia, Epigastric distress, Nausea, Vomiting, etc.
Urinary	Urination impaired, etc.

15. OTHER PRECAUTIONS

15.1 Information Based on Clinical Use

Eczema or dermatitis may be aggravated.

20. PRECAUTIONS FOR HANDLING

20.1 To maintain the quality of the product, avoid moisture as much as possible and store in a cool place, away from direct sunlight.

20.2 Avoid moisture, especially after opening, and handle with care.

20.3 Since this product is made from crude drugs, the color of the product may vary.

22. PACKAGING

(KB-1)

3.75 g \times 28 packets [sachets]

3.75 g \times 168 packets [sachets]

(EK-1)

2.5 g \times 42 packets [sachets]

2.5 g \times 294 packets [sachets]

500 g [bottle]

26. MARKETING AUTHORIZATION HOLDER, etc.

26.1 Manufactured and Distributed by:

Kracie, Ltd.

20-20, Kaigan 3-Chome, Minato-ku, Tokyo 108-8080, Japan

26.2 Distributed by:

Kracie Pharmaceutical, Ltd.

20-20, Kaigan 3-Chome, Minato-ku, Tokyo 108-8080, Japan